
Oxbow Lake

Aquatic Plant Management Plan – 3rd edition

Prepared for:

Presque Isle Town Lake Committee

Contact: Nick Williams

P.O. Box 37

Presque Isle, Wisconsin 54557

Prepared by:

White Water Associates, Inc.

Dean Premo Ph.D., and Carole Gabbard, M.S.

429 River Lane, P.O. Box 27

Amasa, Michigan 49903

Phone: (906) 822-7889; E-mail: dean.premo@white-water-associates.com

Cite as: Premo, Dean and Carole Gabbard. 2025. *Oxbow Lake Aquatic Plant Management Plan*. 3rd edition. White Water Associates, Inc.



Date: April 3, 2026

Presque Isle Wilderness Waters Program

Aquatic Plant Management Plan – Oxbow Lake

This plan is a product of a WDNR Lake Planning Grant awarded to:

Presque Isle Town Lakes Committee

P.O. Box 37

Presque Isle, Wisconsin 54557

Contact: Nick Williams

Phone: (715) 892-7184; E-mail: thewds@milwpc.com

Submitted to:

Wisconsin Department of Natural Resources

Attention: Kevin Gauthier, Sr., Water Resource Management Specialist

8770 Hwy J

Woodruff, WI 54568

Phone: (715) 365-5211 ext. 214; Email: Kevin.GauthierSr@wisconsin.gov

Prepared by:

White Water Associates, Inc.

Dean Premo Ph.D., and Carole Gabbard, M.S.

429 River Lane, P.O. Box 27

Amasa, Michigan 49903

Phone: (906) 822-7889; E-mail: dean.premo@white-water-associates.com



Table of Contents

| | |
|--|----|
| Chapter 1. Introduction..... | 1 |
| Chapter 2. Study area..... | 4 |
| Chapter 3. Purpose and goal statements..... | 8 |
| Chapter 4. Information and Analysis..... | 9 |
| Section 1. Watershed..... | 9 |
| Section 2. Aquatic plant management history..... | 13 |
| Section 3. Basic knowledge on aquatic plants..... | 13 |
| Section 4. Oxbow Lake aquatic plant community..... | 14 |
| Part 1. Oxbow Lake aquatic plants in 2024..... | 15 |
| Part 2. Oxbow Lake aquatic plant community history..... | 27 |
| Part 3. Oxbow Lake aquatic plant community highlights..... | 35 |
| Section 5. Fish community..... | 36 |
| Section 6. Water quality and trophic status..... | 36 |
| Section 7. Water use..... | 36 |
| Section 8. Riparian area..... | 36 |
| Section 9. Wildlife..... | 37 |
| Section 10. Stakeholders..... | 37 |
| Chapter 5. Recommendation, actions, and objectives..... | 38 |
| Chapter 6. Contingency plan for AIS..... | 43 |
| Appendix 1. Literature cited..... | 46 |
| Appendix 2. Documents..... | 48 |
| Appendix 3. Aquatic Invasive Species Report..... | 61 |



Table of Exhibits

| | |
|--|----|
| Map 1. Presque Isle Township lakes on which point-intercept aquatic plant surveys have been conducted. | 4 |
| Map 2. Aerial photo of Oxbow Lake. | 5 |
| Map 3. Topographic map of Oxbow Lake. | 7 |
| Map 4. Watersheds of Presque Isle Township Area. | 10 |
| Map 5. Digital elevation model of Presque Isle Township area. | 11 |
| Map 6. 2024 substrate distribution in North Part of Oxbow Lake. | 18 |
| Map 7. 2024 substrate distribution in South Part of Oxbow Lake. | 19 |
| Map 8. North Part of Oxbow Lake total rake fullness per PI point in 2024. | 20 |
| Map 9. South Part of Oxbow Lake total rake fullness per PI point in 2024. | 21 |
| Map 10. North Part of Oxbow Lake number of aquatic plant species documented per PI point in 2024. | 23 |
| Map 11. South Part of Oxbow Lake number of aquatic plant species documented per PI point in 2024 | 24 |
| Map 12. 2014, 2018, and 2024 total rake fullness distribution in North Part of Oxbow Lake | 30 |
| Map 13. 2014, 2018, and 2024 total rake fullness distribution in South Part of Oxbow Lake. | 31 |
| | |
| Table 1. Water body parameters | 6 |
| Table 2. Cover Types and Soil Groups of the Oxbow Lake Watershed..... | 12 |
| Table 3. Summary statistics for the 2024 PI aquatic plant surveys for Oxbow Lake..... | 16 |
| Table 4. Summary statistics for the 2007, 2014, 2018 and 2024 PI aquatic plant surveys for Oxbow Lake..... | 29 |



| | |
|--|----|
| Figure 1. Percentages of PI points with various number of species per point (Note: "N" is the total number of species found at a point)..... | 22 |
| Figure 2. Oxbow Lake aquatic plants relative frequency in 2024. | 25 |
| Figure 3. Oxbow Lake relative frequencies for the plants collected during the 2007, 2014, 2018 and 2024 PI aquatic plant surveys. | 33 |
| Figure 4. Oxbow Lake frequencies of occurrences for the plants collected during the 2014, 2018 and 2024 PI aquatic plant surveys. | 34 |
| Figure 5. Aquatic Invasive Plant Species Rapid Response Flow Chart | 45 |
| | |
| Appendix 2 - Table 1. Description of the aquatic plant community statistical terms | 49 |
| Appendix 2 - Table 2. Plant species recorded and statistics for the 2024 Oxbow Lake aquatic plant survey. | 51 |
| | |
| Appendix 2 - Map 1. North Part of Oxbow Lake total rake fullness distribution during the 2014 aquatic plant survey. | 55 |
| Appendix 2 - Map 2. North Part of Oxbow Lake total rake fullness distribution during the 2018 aquatic plant survey. | 56 |
| Appendix 2 - Map 3. North Part of Oxbow Lake total rake fullness distribution during the 2024 aquatic plant survey. | 57 |
| Appendix 2 - Map 4. South Part of Oxbow Lake total rake fullness distribution during the 2014 aquatic plant survey. | 58 |
| Appendix 2 - Map 5. South Part of Oxbow Lake total rake fullness distribution during the 2018 aquatic plant survey. | 59 |
| Appendix 2 - Map 6. South Part of Oxbow Lake total rake fullness distribution during the 2024 aquatic plant survey. | 60 |



Chapter 1. Introduction

The *Presque Isle Wilderness Waters Program* results from the efforts of the Presque Isle Town Lakes Committee, an organization that has been active since 2005. The Lakes Committee views stewardship of lakes as an ongoing endeavor that is integrated, coordinated, and administered by the Lakes Committee. This broader perspective accommodates the appropriate range of geographic scales from which to approach lake stewardship: a discrete “lake specific” focus that goes hand-in-hand with waterscape-wide awareness.

This aquatic plant management plan addresses Oxbow Lake. Despite this specificity, it maintains the waterscape perspective crucial to effective lake stewardship. This is especially important when it comes to preventing introduction and establishment of aquatic invasive species (AIS). The closely related *Wilderness Waters Adaptive Management Plan* (Stine et al., 2022) provides additional overarching waterscape level examination that allows greater opportunity and efficiency in water resource management and education.

A 2024 systematic survey of aquatic plants using the Wisconsin Department of Natural Resources (WDNR) “point-intercept” method was an important underpinning of this aquatic plant management plan. An analysis of the plant data along with water quality and other lake information allowed the preparation of the plan.

Aquatic plants rarely get the respect they merit, although this is slowly changing. We still call an aquatic plant bed a “weed bed.” Many aquatic plants have “weed” in their names (e.g., duckweed, pondweed, or musky weed). Likely this term was borrowed from “seaweed” and not intended as derogatory, but in today’s use, “weed” connotes an unwanted, aggressively growing plant. Such is not the case for the vast majority of aquatic plants. In fact, aquatic plants are a vital part of a lake ecosystem, recycling nutrients, providing vertical and horizontal structure, and creating habitat for animal life. Invertebrates, including crustaceans and insects, live on or within this “aquatic forest.” Fish find food and shelter within aquatic plant beds. Waterfowl eat parts of plants directly as well as feed on invertebrates associated with the plants. Muskrats eat aquatic plants and particularly love cattails and bulrushes. Otter and mink hunt invertebrates and small vertebrates within the shelter of submergent and emergent beds. In shallow water, great blue herons find fishes among the plants.

In lakes that receive an excess of nutrients (particularly from fertilizers or leaking septic tanks), plant growth can become too dense or dominated by only a few species. As these abundant plants die, their decomposition can depress dissolved oxygen levels and diminish suitability for fish. Algae can respond rapidly to nutrient influxes and create nuisance conditions. These phenomena can cause people to view all aquatic plants in a negative light.



On another negative front, non-native plant species, transported on boats and trailers or dumped from home aquariums, private ponds or water gardens may proliferate in a water body and negatively influence the community of native species. Eurasian water-milfoil (*Myriophyllum spicatum*) is one of the invasive plant species capable of this kind of population boom. Fortunately, this kind of rampant growth of aquatic invasive plants does not always occur. On occasion, even a native plant species can exhibit excessive growth and results in a population that is viewed by some as a recreational nuisance. The native Southern Naiad (*Najas guadalupensis*) has exhibited this kind of behavior in some northern Wisconsin lakes.

For most lakes, native aquatic plants are an overwhelmingly positive attribute, greatly enhancing the aesthetics of the lake and providing good opportunities for fishing, boating, swimming, snorkeling, sight-seeing, and hunting. In some water bodies even the presence of an aquatic invasive plant species is not a significantly negative phenomenon.

When it comes to aquatic plant management, it is useful to heed the mantra of the medical profession: “First, do no harm.” It is both a social and scientific convention that aquatic plant management is more effective and beneficial when a lake is considered as an entire and integrated ecosystem. Actions taken to curtail a specific plant population (for example, herbicide use to treat Eurasian watermilfoil) will invariably impact other desirable native species. Rare plants, important habitats, or culturally significant plants (such as wild rice) should always be given careful consideration and protection.

Anyone involved in aquatic plant management should be aware that a permit may be required to remove, add, or control aquatic plants. In addition, anyone using Wisconsin’s lakes must comply with the “Boat Launch Law” that addresses transport of aquatic plants on boat trailers and other equipment. A good review of the laws, permits, and regulations that affect management and behavior surrounding aquatic plants can be found in the WDNR guidelines called *Aquatic Plant Management in Wisconsin*.

In preparing this plan, we followed guidelines in *Aquatic Plant Management in Wisconsin*. The resulting plan is an adaptive plan (Walters, 1986). Simply put, it will be modified as new information becomes available. The WDNR Guidance document outlines three objectives that may influence preparation of an aquatic plant management plan. Currently, the motivation for this plan lies in the first two objectives:

- **Protection** - preventing the introduction of nuisance or invasive species into waters where these plants are not currently present;

- **Maintenance** - continuing the patterns of recreational use that have developed historically on and around a lake; and



• **Rehabilitation** - controlling an imbalance in the aquatic plant community leading to the dominance of a few plant species, frequently associated with the introduction of invasive non-native species.

During projects with the WDNR Planning Grant Program and through past efforts, the Town Lakes Committee has followed the seven-step plan outlined in the Guidance Document for developing an aquatic plant management plan:

1. Goal setting – Getting the effort organized, identifying problems to be addressed, and agreeing on the goals;
2. Inventory – Collecting baseline information to define the past and existing conditions;
3. Analysis – Synthesizing the information, quantifying and comparing the current conditions to desired conditions, researching opportunities and constraints, and setting directions to achieving the goals;
4. Alternatives – Listing possible management alternatives and evaluating their strengths, weaknesses and general feasibility;
5. Recommendations – Prioritizing and selecting preferred management options, setting objectives, drafting the plan;
6. Implementation – Formally adopting the plan, lining up funding, and scheduling activities for taking action to achieve the goals;
7. Monitor & Modify – Developing a mechanism for tracking activities and adjusting the plan as it evolves.


Besides this introductory chapter, this plan is organized in six Chapters. The study area is described in Chapter 2. Chapter 3 states the purpose and goals for the plan. Chapter 4 presents an inventory and analysis of information that pertain to the plan including the results of the aquatic plant survey. Chapter 5 provides recommendations that support the overall goals and establish the stewardship component of the plan. This chapter also presents actions and objectives for implementing the plan. Finally, Chapter 6 describe the contingency plan for aquatic invasive species. Three appendices complete this document. Appendix 1 contains literature cited, Appendix 2 contains some tables and figures of the aquatic plant survey results, Appendix 3 contains the 2024 Aquatic Invasive Species Report for Oxbow Lake.



Chapter 2. Study area

“Almost an island” is the literal translation of the French phrase “Presque Isle.” Early French missionaries, perhaps disoriented by the preponderance of water in this north central Wisconsin landscape applied the name, “Presque Isle” to describe an area where the water seemed to dominate the land. The French visitors and Native Americans certainly recognized this landscape as special. Modern ecologists and recreationists share this view. The region that includes the Township of Presque Isle, Wisconsin is an ecological landscape marvelously rich in surface waters. Aerial photography reveals a concentration of lakes and streams that is unique in North America. Presque Isle Township has eighty-four lakes. The Presque Isle area could as easily be termed a “waterscape” as a “landscape.”

Presque Isle Township is one of the northern-most townships in Vilas County, Wisconsin. Presque Isle Township’s northern border is shared with the State of Michigan. In fact, some of the Presque Township lakes lie on the state border. The location of the subject of this APM Plan (Oxbow Lake) is shown in Map 1 along with other lakes in Presque Isle Township that have had point-intercept aquatic plant surveys conducted.



Map 1. Presque Isle Township lakes on which point-intercept aquatic plant surveys have been conducted.





Map 2. Aerial photo of Oxbow Lake.



Descriptive parameters for Oxbow Lake are in Table 1. It is a drainage lake of about 523 acres and maximum depth of 44 feet. Oxbow Lake has a Shoreline Development Index (SDI) of 4.2. The SDI is a quantitative expression derived from the shape of the lake defined as the ratio of the shoreline length to the length of the circumference of a circle of the same area as the lake. A perfectly round lake has an index of 1. Increasing irregularity of a lake’s shoreline in the form of bays and projections of the shore is results in SDIs greater than 1. For example, fjord lakes with extremely irregular shorelines sometimes have SDI’s exceeding 5. A higher shoreline development index indicates that a lake has relatively more productive littoral zone habitat. Oxbow Lake’s shoreline in one of the highest SDI we have observed on inland lakes.

Table 1. Water body parameters

| | |
|-------------------------------------|--|
| Water Body Name | Oxbow |
| County | Vilas |
| Township/Range/Section | T43N-R06E-S1; T43N-R07E-S6; T44N-R06E-S36; T44N-R07E-S31 |
| Water Body Identification Code | 2954800 |
| Lake Type | Drainage |
| Surface Area (acres) | 523 |
| Maximum Depth (feet) | 44 |
| Maximum Length (miles) | 1.5 |
| Maximum Width (miles) | 1.25 |
| Shoreline Length (miles) | 13.5 |
| Shoreline Development Index | 4.2 |
| Total Number of Piers (2020 aerial) | 98 |
| Number of Piers / Mile of Shoreline | 7.3 |
| Total Number of Homes (2020 aerial) | 151 |
| Number of Homes / Mile of Shoreline | 11.2 |



Oxbow Lake has one public access site. A total of 98 piers on the shoreline of Oxbow Lake was observed from recent aerial photography or about 7.3 piers per mile of shoreline. The riparian area is of high quality and undisturbed in many areas. It consists of both upland and wetland areas (Map 3).



Map 3. Topographic map of Oxbow Lake.



Chapter 3. Purpose and goal statements

This plan approaches aquatic plant management with a healthy dose of humility. We do not always understand the causes of environmental phenomena or the effects of our actions to manage the environment. With that thought in mind, we have crafted a statement of purpose and goals for this plan:

Oxbow Lake has a native aquatic plant community that has been documented with point-intercept aquatic plant surveys conducted in 2007, 2014, and 2018. This plant community is essential to, and part of, a high-quality aquatic ecosystem that benefits the human community with its recreational and aesthetic features. The purpose of this aquatic plant management plan is to maintain the aquatic plant community in its present high-quality state.

Supporting this purpose, the goals of this aquatic plant management plan are:

- (1) Monitor and protect the native aquatic plant community;*
- (2) Monitor for AIS and prevent establishment of new non-native biota;*
- (3) Consider and evaluate the efficacy of active aquatic plant management; and*
- (4) Educate riparian owners and lake users on preventing AIS introduction, reducing nutrient inputs that can alter the plant community, minimizing physical removal of native riparian and littoral zone plants, and living with a lake whose natural healthy state includes aquatic plants.*

The purpose and goals are the foundation for the aquatic plant management plan presented in this document. They inform the objectives and actions outlined in Chapter 5 and are the principal motivation of Oxbow Lake stewards.




Chapter 4. Information and Analysis

Our efforts in the Wilderness Waters Program have compiled information about historical and current conditions of the Oxbow Lake ecosystem and its surrounding watershed. Of particular importance to this aquatic plant management plan is the aquatic plant survey that was conducted using the *WDNR Protocol for Aquatic Plant Survey, Collecting, Mapping, Preserving, and Data Entry* (Hauxwell et al., 2010). The results of this comprehensive “point-intercept” survey along with relevant components of other information are presented in this chapter under nine respective subheadings: watershed, aquatic plant management history, aquatic plant community description, fish community, water quality and trophic status, water use, riparian area, wildlife, and stakeholders.

Section 1. Watershed

The Presque Isle Township waterscape sits on a large-scale watershed divide. Some of the water drains north through the Presque Isle River system and eventually enters Lake Superior. Some of the water drains into the Flambeau River system and on to the Mississippi River. In fact, there are two federal hydrologic sub-basins (designated by 8-digit HUC codes) that include Presque Isle Township. The Black-Presque Isle Rivers sub-basin (HUC#04020101) drains north to Lake Superior and the Flambeau River sub-basin (HUC#0705002) drains southwesterly to the Mississippi River. The Black-Presque Isle Rivers sub-basin contains two federal hydrologic sub-watersheds within Presque Isle Township: the South Branch Presque Isle River sub-watershed (HUC#040201010303) and the Pomeroy Creek-East Branch Presque Isle River sub-watershed (HUC#040201010301). The Flambeau River sub-basin contains one sub-watershed within Presque Isle Township: the Rice Creek sub-watershed (HUC#07050020103). Map 5 illustrates these watersheds and the watersheds of the water bodies subject to the Wilderness Waters Program studies. Oxbow Lake is contained within the South Branch Presque Isle River sub-watershed (Map 4).






Map 4. Watersheds of Presque Isle Township Area.

Description: Portions of 3 federal hydrologic sub-watersheds are illustrated: (1) S.Br. Presque Isle River (bounded by yellow), (2) Pomeroy Cr-E.Br. Presque Isle River (bounded by blue and yellow), and (3) Rice Cr. (bounded by green, yellow, and blue). Also shown are the smaller watersheds of individual water bodies subject to the Wilderness Waters studies (bounded by red and, in the case of Papoose Lake, bounded by white). White dots show outlet points for the smaller watersheds.



The elevation in Presque Isle Township ranges from around 1,550 feet above sea level to 1,750 feet above sea level. A digital elevation model is provided as Map 5 and shows the relative elevations for the area with orange areas of the landscape being the highest elevations and greens and blues being the lowest elevations.



Map 5. Digital elevation model of Presque Isle Township area.

The watershed (drainage basin) is all of the land and water areas that drain toward a particular river or lake. A water body is greatly influenced by its watershed. Watershed size, topography, geology, land use, soil fertility and erodibility, and vegetation are all factors that influence water quality. The Oxbow Lake watershed is about 5,925 acres. The cover types in the watershed are presented in Table 2. Forest and surface water comprise the largest components. Soil group B is most prevalent, followed by group D, while groups A and C make up around 20%. Soil group A has the highest infiltration capacity, and the lowest runoff potential. Conversely, soil group D has the lowest infiltration capacity, and the highest runoff potential. The watershed to lake area ratio is 11:1. Water quality often decreases with an increasing ratio of watershed area to lake area. As the watershed to lake area increases there are more sources of runoff. In larger



watersheds, runoff water can leach more minerals and nutrients and carry them to the lake. Runoff to a lake (such as after a rainstorm or snowmelt) differs greatly among land uses. Forest cover is the most protective as it exports much less soil (through erosion) and nutrients (such as phosphorus and nitrogen) to the lake than agricultural or urban land use.

Table 2. Cover Types and Soil Groups of the Oxbow Lake Watershed.

| Cover Type | | Acres | Percent |
|--------------------------|--------|---------|---|
| Agriculture | | 0 | 0 |
| Commercial | | 0 | 0 |
| Forest | | 3445.9 | 58.2 |
| Grass/Pasture | | 4.2 | 0.1 |
| High-density Residential | | 0 | 0 |
| Low-density Residential | | 336.6 | 5.7 |
| Water | | 2136.7 | 36.1 |
| Total | | 5923.4 | 100.0 |
| Soil Group | Acres | Percent | Hydrologic Soil Groups - Soils are classified by the Natural Resource Conservation Service into four Hydrologic Soil Groups* based on the soil's runoff potential. The four Hydrologic Soils Groups are A, B, C and D. Where A has the smallest runoff potential and D the greatest. |
| A | 893.5 | 15.1 | Group A is sand, loamy sand or sandy loam types of soils. It has low runoff potential and high infiltration rates even when thoroughly wetted. They consist chiefly of deep, well to excessively drained sands or gravels and have a high rate of water transmission. |
| B | 2680.8 | 45.3 | Group B is silt loam or loam. It has a moderate infiltration rate when thoroughly wetted and consists chiefly or moderately deep to deep, moderately well to well drained soils with moderately fine to moderately coarse textures. |
| C | 318.8 | 5.4 | Group C soils are sandy clay loam. They have low infiltration rates when thoroughly wetted and consist chiefly of soils with a layer that impedes downward movement of water and soils with moderately fine to fine structure. |
| D | 2030.2 | 34.3 | Group D soils are clay loam, silty clay loam, sandy clay, silty clay or clay. This soil has the highest runoff potential. They have very low infiltration rates when thoroughly wetted and consist chiefly of clay soils with high swelling potential, soils with a permanent high water table, soils with a claypan or clay layer at or near the surface and shallow soils over nearly impervious material. |



Section 2. Aquatic plant management history

As far as we can determine, no systematic or large-scale plant management activity has ever taken place in Oxbow Lake. Over the years, no particular nuisance issues have demanded control action. Systematic aquatic plant surveys have been conducted on Oxbow Lake in 2007, 2014, 2018 and 2024. Findings from the 2007, 2014, and 2018 surveys are discussed and compared to the most recent (2024) survey in Part 2.

Section 3. Basic knowledge on aquatic plants

Why do lakes need aquatic plants? In many ways, they are underwater forests. Aquatic plants provide vertical and horizontal structure in the lake just like the many forms and variety of trees do in a forest. Imagine how diminished a forest's biodiversity becomes in the advent of a clear-cut or a conversion from a hardwood forest to a red pine plantation. Similarly, a lake's biodiversity in large part depends on a diversity of plants.

Aquatic plants are beneficial in many ways. Areas with plants produce more food for fish in the form of insect larvae, snails, and other invertebrates. Aquatic vegetation offers fish shelter and spawning habitat. Many submerged plants provide food for waterfowl and habitat for insects on which some waterfowl feed. Aquatic plants further benefit lakes by producing oxygen and absorbing nutrients (phosphorus and nitrogen) from runoff. Aquatic plants also protect shorelines and lake bottoms by dampening wave action and stabilizing sediments.

The distribution of plants within a lake is generally limited by light availability. Light availability at the lake bottom is affected by water clarity. Aquatic biologists often estimate the depth to which rooted aquatic plants can exist as about two times the average Secchi clarity depth. For example, if the average Secchi depth is eight feet, then it is fairly accurate to estimate that rooted plants might exist in water as deep as sixteen feet. At depths greater than that (in this example), light is insufficient for rooted plants to grow. In addition to available light, the type of substrate influences the distribution of rooted aquatic plants. Plants are more likely to be found in muddy or soft sediments containing organic matter, and less likely to occur where the substrate is sand, gravel, or rock. Finally, water chemistry influences which plants are found in a body of water. Some species prefer alkaline lakes and some prefer more acidic lakes. The presence of nutrients like phosphorous and nitrogen also influence plant community composition. Water



quality information for Oxbow Lake can be found in the DNR Water Explorer tool ([WEx WDNR - Oxbow Lake \(Vilas County\)](#))

As mentioned earlier, non-native invasive plant species can reach high densities and wide distribution within a lake. This can diminish the native plant community and the related habitat. At times, even a native plant species can reach high population levels and interfere with certain kinds of human recreation. Cases such as these often elicit calls for some kind of plant management. It should be noted, however, that altering aquatic plant communities through hand-pulling, mechanical harvest, herbicides, or other means is expensive (in time and/or money) and by no means permanent. Long-term outcomes of these manipulations are difficult to predict and collateral damage to non-target plant species can be significant. For example, the efficacy and widespread effects of new chemicals developed for plant management are poorly understood. Safety evaluation of these herbicides often does not include their breakdown products that will reside in the lake environment for unknown duration. Management treatments can even exacerbate the perceived problem or create other management challenges. Because of these uncertainties, permits are typically required in cases of aquatic plant management.

Section 4. Oxbow Lake aquatic plant community

Aquatic plant surveys have been conducted on Oxbow Lake by aquatic plant specialists in 2007, 2014, 2018, and 2024. In each of these surveys, WDNR point-intercept protocol and methodology¹ was followed. This systematic survey assessed the plant species composition on a grid of 843 points distributed evenly over the lake. Using latitude-longitude coordinates and a handheld GPS unit, scientists navigated to the points and used a rake mounted on a pole or rope to sample plants. These were identified, recorded, and put into a dedicated spreadsheet for storage and data analysis. During the preparation of this plan, the data from the four PI surveys (2007, 2014, 2018, and 2024) have been analyzed and interpreted to evaluate the composition of the plant community (species present and abundance) and how the plant community has changed over time.

¹ [Wisconsin Standard Method to Collect Aquatic Plant Data](#)



The aquatic plant data for Oxbow Lake along with data from many Wisconsin Lakes, can be explored in the Wisconsin DNR Aquatic Plant Explorer Tool at [Aquatic Plant Explorer](#) .

Because Oxbow Lake has been surveyed four times, we are able to study the plant community over time. Changes in a lake environment might manifest as loss of species, change in species abundance or distribution, difference in the relative composition of various plant life forms (emergent, floating leaf, or submergent plants), and/or appearance of an AIS or change in its population size. Monitoring can track changes and provide valuable insight on which to base management decisions. In the remainder of this section (Part 1) we provide a report of the findings of the 2024 point-intercept aquatic plant survey and summarize and compare findings of the 2007, 2014, 2018, and 2024 surveys in Part 2.

Part 1. Oxbow Lake aquatic plants in 2024

In 2024, a total of 562 of the 843 point-intercept (PI) points were visited. Of these, 384 points were shallower than the maximum depth of plants. The maximum depth of plant colonization was determined during the 2024 survey to be 13.5 feet (Table 3). Areas in Oxbow Lake at this depth and shallower can be considered the littoral zone. Deeper areas in the lake do not have sufficient light for rooted vegetation to establish. Rooted vegetation was actually found at 225 of the 384 sample points within the littoral zone (58.59% of sites within the littoral zone). This indicates that although availability of sufficient light limits the distribution of plants, it is not the only habitat factor involved. Substrate is another feature that influences plant distribution (e.g., soft substrate often harbors more plants than hard substrate). Maps 6 and 7 presents the substrates encountered during the aquatic plant survey (muck, sand, or rock). The substrate types were evenly represented with 54.7% for muck followed by sand with 24.6% and 20.8% of the substrate is rock.

Plant density is estimated by a “rake fullness” metric (scores are assessed as 1, 2, or 3 with 3 being the highest possible density). The average rake fullness for Oxbow Lake in 2024 was 1.38 (Table 3). These densities (considering all species) are displayed for each sampling point on Maps 8 and 9.




Table 3. Summary statistics for the 2024 PI aquatic plant surveys for Oxbow Lake.

| General Statistics: | 2024 |
|---|-------------|
| Total number of sites on grid - Total number of sites on original grid (not necessarily visited) | 843 |
| Total number of sites visited - Total number of sites where the boat stopped to sample. | 562 |
| Total number of sites with vegetation - Total sites with at least one plant was found | 225 |
| Total number of sites shallower than maximum depth of plants - Number of sites where depth was less than or equal to the maximum depth where plants were found. This value is used for frequency of occurrence at sites shallower than maximum depth of plants. | 384 |
| Frequency of occurrence at sites shallower than maximum depth of plants - Not every area of the lake bottom that is shallower than the maximum depth of plants actually has plants living there. Other things (like substrate) can influence where a plant can root. Frequency of occurrence at sites shallower than max depth of plants documents the proportion of sites where rooted plants actually occur within the littoral zone | 58.59 |
| Simpson Diversity Index - An index that accounts for richness (number of species) and evenness of distribution in estimating diversity. It is based on a plant's relative frequency. Values range from 0 to 1 with higher values indicating greater diversity. | 0.92 |
| Maximum depth of plants (ft) - Since rooted aquatic plants require sunlight at the lake bottom in order to root and grow, water depth and transparency limit where plants can establish. By repeated sampling, we determine the max depth at which plants grow. Plants are absent beyond that depth. The area of the lake where plants can grow is the littoral zone. | 13.5 |
| Number of sites sampled using rake on Rope | 8 |
| Number of sites sampled using rake on Pole | 464 |
| Average number of all species per site (shallower than max depth) - This is the average number of species found at PI points shallower than max depth of rooted plants. Lakes with diverse plant communities tend to have high average number of species per site. | 1.41 |
| Average number of all species per site (veg. sites only) - This is the average number of plant species documented at PI sampling points where plants were collected. Lakes with diverse plant communities tend to have high average number of species at each site | 2.40 |




| General Statistics: | 2024 |
|--|-------------|
| Average number of native species per site (shallower than max depth) - This metric is the average number of native plant species documented at PI sampling points. Lakes with diverse plant communities tend to have high average number of plant species at each site. | 1.41 |
| Average number of native species per site (veg. sites only) - This metric is the average number of native plant species documented at PI points where plants were collected. Lakes with diverse plant communities tend to have high average number of plant species per site. | 2.40 |
| Species Richness - This metric is a simple measure of diversity and is a count of the number of species. In the case of a PI survey, it is the number of species collected on the rake for PI points in a given area (an entire lake or a plot). The species richness of a lake would mean the total number of plant species in the entire lake. More species equals higher diversity | 38 |
| Species Richness (including visuals) - This metric includes the count of the number of plant species collected on the rake for the point-intercept sampling points in a given area and those observed (but not actually collected with the rake) within 6 feet of the sampling points. | 42 |
| FQI – A metric to evaluate the closeness of the flora of an area to that of undisturbed areas. | 45.70 |
| Average Rake Fullness - Rake fullness is a rating given to each rake pull on a PI survey that indicates the amount of plant material for each species as well as combined species. There are four ranks: 0 (not present), 1, 2, & 3 (3 is the highest amount of plant material on the rake). Average rake fullness is the mean value for all of the PI points. | 1.38 |






Map 6. 2024 substrate distribution in North Part of Oxbow Lake.






Map 7. 2024 substrate distribution in South Part of Oxbow Lake.





Map 8. North Part of Oxbow Lake total rake fullness per PI point in 2024.





Map 9. South Part of Oxbow Lake total rake fullness per PI point in 2024.



Species richness is an ecological term that refers to the total number of species in a community (Table 3). Forty-two species of aquatic plants were documented on Oxbow Lake in 2024. Of these, thirty-eight were collected at point-intercept sampling points and the others were observed from the boat while traveling between sampling points. Appendix 2 - Table 2 provides a list of the species encountered, including common and scientific names along with summarizing statistics. The average number of species encountered at the vegetated points was 2.40. The number of species at these vegetated points ranges from one to nine. Figure 1 shows the percentages of vegetated PI points with 1-9 species. Sites with only one, two and three species comprised 80.9% of the sites. The actual number of species encountered at each of the vegetated points is graphically displayed on Maps 10 and 11.






Figure 1. Percentages of PI points with various number of species per point (Note: "N" is the total number of species found at a point).





Map 10. North Part of Oxbow Lake number of aquatic plant species documented per PI point in 2024.





Map 11. South Part of Oxbow Lake number of aquatic plant species documented per PI point in 2024



Appendix 2 - Table 2 provides information about occurrences of the plant species recorded in Oxbow Lake in 2024. Several metrics are provided, including total number of points at which each species was found and frequency of occurrence at points within the littoral zone. This frequency metric is also standardized as a “relative frequency” (also shown in Appendix 2 - Table 2) by dividing the frequency of occurrence for a given species by the sum of frequency of occurrence for all plants and multiplying by 100 to form a percentage. The resulting relative frequencies for all species total 100%. The relative frequencies for the plant species collected with a rake are graphically displayed in alphabetic order on Figure 2. This display shows that *Najas flexilis* (slender naiad) has the highest relative frequency (17.4%) followed by *Brasenia schreberi* (Watershield) with 10% of relative frequency. *Potamogeton amplifolius* (Large-leaf pondweed), *Chara sp.* (Muskgrasses), *Potamogeton berchtoldii* (Berchtold's pondweed) and *Najas gracillima* (Northern naiad) had similar relative frequency in 2024 ranging from 8.9 to 7.6% of the vegetated points. The next four species had lower relative frequencies ranging from 3.3% to 5.2%. A great number of species (28 total) have individually a relative frequency under 3% but comprise a total 23.1% of relative frequency.




Figure 2. Oxbow Lake aquatic plants relative frequency in 2024.



As discussed earlier, “species richness” is the term given to the total number of species in a given area. For example, the total number of plant species in a lake would be its plant species richness. Generally speaking, a high species richness means high biodiversity and this is considered a healthy and desirable condition in an ecosystem. But species richness doesn’t tell the whole story. As an example, consider the plant communities of two hypothetical ponds each with 1,000 individual plants representing ten plant species (in other words, species richness is 10 in both ponds). In the first pond each of the ten species populations is comprised of 100 individuals. In the second pond, Species #1 has a population of 991 individuals and each of the other nine species is represented by one individual plant. Intuitively, we would say that first pond is more diverse because there is more “even” distribution of individual species. The “Simpson Diversity Index” takes into account both richness and evenness in estimating diversity. It is based on a plant’s relative frequency in a lake. The closer the Simpson Diversity Index is to 1, the more diverse the plant community. The Simpson Diversity Index for Oxbow Lake aquatic plants was 0.92 in 2024 (Table 3) which indicates a diverse and healthy aquatic plant community.

Another measure of diversity and quality is the Floristic Quality Index (FQI). Floristic quality is an assessment metric designed to evaluate the closeness that the flora of an area is to that of undisturbed conditions (Nichols 1999). Among other applications, it forms a standardized metric that can be used to compare the quality of different lakes (or different locations within a single lake) and monitor long-term changes in a lake’s plant community (an indicator of lake health). The FQI for a lake is determined by using the average coefficient of conservatism times the square root of the number of native plant species present in the lake. Knowledgeable botanists have assigned to each native aquatic plant a coefficient of conservatism representing the probability that a plant is likely to occur in pristine environments (relatively unaltered from presettlement conditions). The coefficients range from 0 to 10, with 10 being assigned to those species most sensitive to disturbance. As more environmental disturbance occurs, the less conservative species become more prevalent.

Nichols (1999) analyzed aquatic plant community data from 554 Wisconsin Lakes to ascertain geographic characteristics of the FQI metric. This is useful for considering how the Oxbow Lake FQI (45.70 in 2024) compares to other lakes and regions. The statewide medians for number of species and FQI are 13 and 22.2, respectively. Oxbow Lake values are high compared to these statewide values. Nichols (1999) determined that there are four ecoregional-lake types groups in Wisconsin: (1) Northern Lakes and Forests lakes, (2) Northern Lakes and Forests



flowages, (3) North Central Hardwoods and Southeastern Till Plain lakes and flowages, and (4) Driftless Area and Mississippi River Backwater lakes. Oxbow Lake is located in the Northern Lakes and Forests lakes group. Nichols (1999) found species numbers (species richness) for the Northern Lakes and Forests lakes group had a median value of 13. Oxbow Lake data demonstrates higher species richness than this value with 42 documented species. Finally, the Oxbow Lake FQI (45.70) is higher than the median value for the Northern Lakes and Forests lakes group (24.3). These findings also support the contention that the Oxbow Lake aquatic plant community is healthy and diverse.

Part 2. Oxbow Lake aquatic plant community history

a. General aquatic plant community statistics

Aquatic plant surveys were conducted in Oxbow Lake in 2007, 2014, 2018 and 2024 using the systematic “point-intercept” method. The same grid of 843 sampling points was used for these four surveys and the resulting data sets formed the principal underpinning of this aquatic plant management plan. The summary statistics for the four surveys are provided in Table 4. Although Oxbow Lake has been surveyed for aquatic plants four times, the 2007 aquatic plant survey was conducted by another company and the available data is incomplete. The 2014, 2018 and 2024 surveys were all conducted by White Water Associates. With the perspective of comparing the four aquatic plant surveys, the consistency between the 2014, 2018, and 2024 plant communities is high. In contrast, the 2007 data have differences that warrant skepticism regarding their correctness. Three examples support this reasoning: (1) the species richness was low for 2007, but consistently high for the 2014, 2018, and 2024 surveys; (2) *Najas guadalupensis* (southern naiad) was reported in 2007 at 11.99% relative frequency, but was not documented in the 2014, 2018 and 2024 surveys.; and (3) *Elodea canadensis* (common waterweed) was reported in 2007 at 24.18% relative frequency, but in the three subsequent surveys was only found at 0.5%, 0.35%, and 0.93% relative frequencies, respectively. Because of this, the comparison of plant communities in this section, considers the three most recent surveys.

We define the littoral zone by the area where the depth is less than or equal to the maximum depth of rooted plants. The maximum depth of plants can change from year to year due to factors such as water clarity, temperature, and lake level. The maximum depth of plants for Oxbow Lake decreased from 12 feet in 2014 to 10 feet in 2018 and then increased in the 2024 survey to 13.5 feet (see Table 4). The mean maximum depth of plants for the four surveys is 11.8 feet.



Species richness (the total number of species found at the sampling points) varied between surveys but has always been high with values of 39 and 38 species in the two most recent surveys (Table 4). It is also interesting that the average number of species per point-intercept site has increased over time from the 2014 to 2024 surveys (Table 4).

The Simpson Diversity Index, which takes into account both richness and evenness in estimating diversity, was consistent in 2014, 2018 and 2024, meaning that the lake had a consistently diverse aquatic plant community (Table 4).

The Floristic Quality Index resulting from the 2014, 2018 and 2024 surveys were all higher than state and the regional average. The Floristic Quality Index (FQI) was consistently high and increasing over the three surveys (Table 4). This relatively stable FQI supports the contention that the Oxbow Lake plant community is healthy, diverse, and indicative of an environment little disturbed by humans (Table 4).

The average rake fullness of all the plants was recorded for the 2014, 2018 and 2024 point-intercept surveys in Oxbow Lake. These values were low and fairly consistent over the 2014 and 2018 surveys and a slight increase is shown for the 2024 survey (Table 4). In other words, plant biomass is slightly increasing judging from the data of these surveys.

Table 4 contains statistics that compare the aquatic plant communities documented in each of the four point-intercept surveys (note, some statistics were not available for 2007). The total number of sites with vegetation increased from 2014 to 2024. The number of sites with vegetation almost doubled from 2018 (120 sites) to 2024 (225 sites). The distributions of the sites with vegetation found during the 2014, 2018 and 2024 aquatic plant surveys are presented in Maps 12 and 13². The frequency of occurrence at sites shallower than maximum depth of plants increased from the 2014 survey to the 2024 survey (more sites had rooted plants in the recent survey).


² Individual maps are presented in Appendix 2 - Maps 1- 6



Table 4. Summary statistics for the 2007, 2014, 2018 and 2024 PI aquatic plant surveys for Oxbow Lake.


| Summary Statistic | 2007 | 2014 | 2018 | 2024 |
|---|-------------|-------------|-------------|-------------|
| Total number of sites on grid | 843 | 843 | 843 | 843 |
| Total number of sites visited | | 584 | 642 | 562 |
| Total number of sites with vegetation | | 111 | 120 | 225 |
| Total number of sites shallower than maximum depth of plants | | 337 | 273 | 384 |
| Frequency of occurrence at sites shallower than maximum depth of plants | 45.47 | 32.94 | 43.96 | 58.59 |
| Simpson Diversity Index | 0.88 | 0.93 | 0.93 | 0.92 |
| Maximum depth of plants (ft.) | 13 | 12 | 10 | 13.5 |
| Number of sites sampled with rake on rope | | 20 | 16 | 8 |
| Number of sites sampled with rake on pole | | 391 | 341 | 464 |
| Average number of all species per site (shallower than max depth) | 0.92 | 0.58 | 1.03 | 1.41 |
| Average number of all species per site (vegetated sites only) | 2.02 | 1.77 | 2.34 | 2.40 |
| Average number of native species per site (shallower than max depth) | 0.92 | 0.58 | 1.03 | 1.41 |
| Average number of native species per site (vegetated sites only) | 2.02 | 1.77 | 2.34 | 2.40 |
| Species Richness | 23 | 29 | 39 | 38 |
| Species Richness (including visuals) | | 37 | 40 | 42 |
| Floristic Quality Index (FQI) | 33.5 | 36.95 | 45.42 | 45.70 |
| Average rake fullness | | 1.08 | 1.04 | 1.38 |





Map 12. 2014, 2018, and 2024 total rake fullness distribution in North Part of Oxbow Lake





Map 13. 2014, 2018, and 2024 total rake fullness distribution in South Part of Oxbow Lake.



b. Analysis of the aquatic plant species composition

The relative frequencies for the plant species collected with a rake for the four point-intercept plant surveys are graphically displayed on Figure 3. Since this graph displays relative frequencies, the total necessarily equals 100%. The graph shows the composition of the plant community in each of the survey years. As one plant increases in relative frequency, others (one or several species) decrease. In the case of Oxbow Lake, we observed no dramatic trends in Figure 3. The graph reflects a very diverse community. The diversity of habitat present in Oxbow Lake is reflected by the diversity of the aquatic plant species.

Figure 4 presents the frequency of occurrence within vegetated areas of aquatic plants between the 2014, 2018 and 2024 surveys. The frequency of occurrences within vegetated areas is the number of sites at which a species was observed divided by the total number of vegetated sites. The graph shows that seven species were first recorded in 2024. One of these species, Vasey's pondweed (*Potamogeton vaseyi*) is a species of special concern in Wisconsin. Figure 4 illustrates a very diverse and dynamic aquatic plant community, species increase and decrease from year to year. Some species are quite common but most are relatively uncommon in the community.






Figure 3. Oxbow Lake relative frequencies for the plants collected during the 2007, 2014, 2018 and 2024 PI aquatic plant surveys.






Figure 4. Oxbow Lake frequencies of occurrences for the plants collected during the 2014, 2018 and 2024 PI aquatic plant surveys.



Part 3. Oxbow Lake aquatic plant community highlights

Lake ecosystems are dynamic - change is the norm rather than the exception. Changes in the ecosystem result from numerous direct and indirect factors and this complexity makes it challenging to determine the causal relationships. Analysis of the three aquatic plant data sets (2014, 2018 and 2024) available for the Oxbow Lake plant community has revealed several important points:

- The total number of sites with vegetation increased from 2014 and 2024, and almost doubled from 2018 to 2024.
- The frequency of occurrence at sites shallower than maximum depth of plants increased over the three most recent surveys (more sites had rooted plants in the recent survey).
- The maximum depth of plants for Oxbow Lake stayed consistent with a mean at 11.8 feet for the three surveys.
- The aquatic plant community diversity has remained consistently high when considering species richness, Simpsons Diversity Index, and Floristic Quality Index. The number of species per point-intercept site has increased over time (2014, 2018, and 2024). These values demonstrate a diverse and healthy aquatic plant community.
- The percent relative frequencies for the species found in Oxbow Lake during the 2024 survey does not show any dramatically dominant species.
- The frequencies of occurrence reflect a very diverse and dynamic aquatic plant community, species increase and decrease from year to year
- Vasey's pondweed (*Potamogeton vaseyi*), species of special concern in Wisconsin, was found for the first time in Oxbow Lake in 2024.
- The average rake fullness showed an increase during the 2024 survey. In other words, plant biomass increased in 2024.
- No aquatic invasive plant species were found in Oxbow Lake.
- One wetland invasive species was confirmed in Oxbow Lake (Appendix 3).



Section 5. Fish community

It was beyond the scope of the current Wilderness Waters project to characterize the fish community and fish habitat of this water body. The WDNR Lake Pages website ([Wisconsin Lakes - Oxbow Lake](#)) indicates that the bottom is comprised of 60% sand, 20% gravel, 15% rock, and 5% muck and that fish species present include musky, panfish, largemouth bass, smallmouth bass, northern pike and walleye.

Section 6. Water quality and trophic status

It was beyond the scope of the current Wilderness Waters project to characterize the water quality of Oxbow Lake. A record of Oxbow Lake water quality data and along with some interpretation can be accessed at the following link: [Water Explorer WDNR - Oxbow Lake](#)

Section 7. Water use

Oxbow Lake has one public access site, and is used by riparian owners and recreationists for a variety of activities. There is a small area of State of Wisconsin Land at the public boat access.

Section 8. Riparian area

Part 1 (Watershed) describes the larger riparian area context of Oxbow Lake. The near shore riparian area can be appreciated by viewing Maps 2 and 3. Our review of recent aerial photography reveals 151 houses on the lake. The lake is lightly developed and has an intact forested riparian zone that extends for hundreds of feet back from the lake. The forest is a mixture of coniferous and deciduous trees and shrubs. This intact riparian area provides numerous important functions and values to the lake. It effectively filters runoff to the lake. It provides excellent habitat for birds and mammals. Trees that fall into the lake from the riparian zone contribute important habitat elements to the lake. Educating riparian owners as to the value of riparian areas is important to the maintenance of these critical areas.

The WDNR drafted in 2016 and formulated a final version in 2020 of a protocol called *Lake Shoreland and Shallows Habitat Monitoring* (WDNR, 2020). It provides a standard methodology for surveying, assessing, and mapping habitat in lakeshore areas, including the Riparian buffer, Bank, and Littoral Zones (WDNR, 2020). This information is useful to local and regional resource managers, community stakeholders, and others interested in protecting and enhancing Wisconsin's



lakes and rivers (WDNR, 2020). The shoreland and shallow water data for Oxbow Lake was recorded in 2018 and the results are available in *Oxbow Lake Aquatic Plant Management Plan 2nd edition* (Premo et al., 2020) or online in WDNR website. The WDNR Lakes and AIS Viewer can be used to see all of the different aspects that were measured in the Shoreland Habitat Survey on a map. This tool can be found here: [WDNR Lake viewer map](#) . Then you will want to click on the Layers tab and the Shoreland Habitat Monitoring box. The WDNR Shoreland Habitat Disturbance Viewer can be found here: [Shoreland Habitat Disturbance Viewer](#) . This link gives both lake wide and property parcel level general statistics on shoreline disturbances and recommendations for improvement.

Section 9. Wildlife

A study of wildlife was beyond the scope of the current study, but would be valuable to study and interpret in future iterations of the plan. This would be especially true of wetland and water-oriented wildlife such as frogs, waterfowl, fish-eating birds, aquatic and semi-aquatic mammals, and invertebrate animals (for example freshwater mussels). Of special importance would be monitoring for the presence of aquatic invasive wildlife species (for example, rusty crayfish, spiny water flea, or zebra mussel) and fish species (for example, rainbow smelt or common carp).

Oxbow Lake is currently designated as a priority navigable water (PNW) (WI Admin. Code, 2020 – the PNW can be found in the WDNR tool online [Wisconsin DNR - Surface Water Data Viewer](#)). Priority Navigable Waters meet any of these standards: navigable waterways, or portions thereof, that are considered ORW/ERW or trout streams; lakes less than 50 acres in size; tributaries and rivers connecting to inland lakes containing naturally-reproducing lake sturgeon populations; waters with self-sustaining walleye populations in ceded territories; waters with self-sustaining musky populations; or perennial tributaries to trout streams (WI Admin. Code, 2020). Oxbow Lake is considered a PNW because it is an ORW and has self-sustaining musky and walleye populations.

Section 10. Stakeholders

At this juncture in the ongoing aquatic plant management planning process, the Town Lakes Committee has represented the Oxbow Lake stakeholders. Additional stakeholders and interested citizens are invited to participate as the plan is refined and updated in order to broaden input, build consensus, and encourage participation in stewardship. No contentious direct plant management actions (for example, harvesting or use of herbicides) are a component of the current plan.



Chapter 5. Recommendation, actions, and objectives

In this chapter we provide recommendations for specific objectives and associated actions to support the APM Plan’s goals stated in Chapter 3 and re-stated here for convenient reference:

- (1) Monitor and protect the native aquatic plant community;*
- (2) Monitor for AIS and prevent establishment of new non-native biota;*
- (3) Consider and evaluate the efficacy of active aquatic plant management; and*
- (4) Educate riparian owners and lake users on preventing AIS introduction, reducing nutrient inputs that can alter the plant community, minimizing physical removal of native riparian and littoral zone plants, and living with a lake whose natural healthy state includes aquatic plants.*

Since Oxbow Lake is a healthy ecosystem, we could simply recommend an alternative of “no action.” In other words, Oxbow Lake continues without any effort or intervention on part of lake stewards. Nevertheless, we consider the “no action” alternative imprudent. Many forces threaten the quality of the lake and Wilderness Waters Program and Town Lakes Committee feels a great responsibility to minimize the threats. We therefore outline in this section a set of actions and related management objectives that will actively engage lake stewards in the process of management.

The actions are presented in tabular form. Each “action” consists of a set of four statements: (1) a declarative “action” statement that specifies the action (2) a statement of the “objective” that the action serves, (3) a “monitoring” statement that specifies the party responsible for carrying out the action and maintaining data, and (4) a “status” statement that suggests a timeline/calendar and indicates status (not yet started, ongoing, or completed).

At this time, we recommend no direct manipulation of plant populations in Oxbow Lake. There are currently no non-native aquatic plants documented for the lake. One wetland invasive plant species was found during the Aquatic Invasive Species Survey (Appendix 3). No native plants exhibit nuisance population size or distribution. Monitoring of the shoreline for wetland invasives should be ongoing and appropriate management employed when needed.



Recommended Actions for the Oxbow Lake APM Plan

Action #1: Formally adopt the Aquatic Plant Management Plan.

Objective: To provide foundation for long-term native plant community conservation and stewardship and to be prepared for response to AIS introductions.

Monitoring: The Lake Association and Town Lakes Committee oversee activity and maintains the plan.

Status: Planned for 2020.

Action #2: Monitor water quality.

Objective: Continue with collection and analysis of water quality parameters to detect trends in parameters such as nutrients, chlorophyll *a*, and water clarity.

Monitoring: The Lake Association or Town Lakes Committee oversees activity and maintains data.

Status: Ongoing.

Action #3: Monitor the lake for aquatic invasive plant species.

Objective: To understand the lake's biotic community, provide for early detection of AIS and continue monitoring any existing populations of AIS.

Monitoring: The Lake Association or Town Lakes Committee oversees activity and maintains data.

Status: Ongoing.

Action #4: Monitor the lake for aquatic invasive animal species.

Objective: To understand the lake's biotic community, provide for early detection of AIS and continue monitoring any existing populations of AIS.

Monitoring: The Lake Association or Town Lakes Committee oversees activity and maintains data.

Status: Ongoing.



Recommended Actions for the Oxbow Lake APM Plan

Action #5: Form an Aquatic Invasive Species Rapid Response Team and interface with the Town Lakes Committee AIS Rapid Response Coordinator.

Objective: To be prepared for AIS discovery and efficient response.

Monitoring: The Lake Association and/or Town Lakes Committee coordinate activity.

Status: Planned for 2020.

Action #6: Conduct quantitative plant survey every five years using WDNR Point-Intercept Methodology.

Objective: To watch for changes in native species diversity, floristic quality, plant abundance, and plant distribution and to check for the occurrence of non-native, invasive plant species.

Monitoring: Town Lakes Committee (Wilderness Waters Program) oversees and maintains data; copies to WDNR.

Status: Anticipated in 2023.

Action #7: Update the APM plan approximately every five years or as needed to reflect new plant information from plant surveys and monitoring.

Objective: To have current information and management science included in the plan.

Monitoring: Lake Association and/or Town Lakes Committee (Wilderness Waters Program) oversees and maintains data; copies to WDNR.

Status: Ongoing.

Action #8: Develop a Citizen Lake Monitoring Network to monitor for invasive species and develop strategies including education and monitoring activities (see <http://www.uwsp.edu/cnr/uwexplakes/clmn> for additional ideas).

Objective: To create a trained volunteer corps to monitor aquatic invasive species and to educate recreational users regarding AIS.

Monitoring: The Lake Association oversees activity and reports instances of possible introductions of AIS.

Status: Anticipated to begin in 2020.



Recommended Actions for the Oxbow Lake APM Plan

Action #9: Become familiar with and recognize the water quality and habitat values of ordinances and requirements on boating, septic, and property development.

Objective: To protect native aquatic plants, water quality, and riparian habitat.

Monitoring: Lake residents and other stakeholders.

Status: Ongoing.

Action #10: Promote adherence to, and enforcement of, the Town of Presque Isle's 200 foot no-wake ordinances (from shoreline and islands).

Objective: To minimize recreational impacts on the aquatic plant community and shoreline habitats, and promote safe boating.

Monitoring: Town Lakes Committee oversees activity and assesses effectiveness.

Status: Ongoing.

Action #11: Create an education plan for the property owners and other stakeholders that will address issues concerning aquatic and riparian plant communities.

Objective: To educate stakeholders about issues and topics that affect the lake's aquatic and riparian plant communities, including topics such as: (1) the importance of the aquatic plant community; (2) no or minimal mechanical removal of plants along the shoreline is desirable and that any plant removal should conform to Wisconsin regulations; (3) the value of a natural shoreline in protecting the aquatic plant community and lake health; (4) nutrient sources to the lake and the role excess nutrients play in degradation of the aquatic plant community; (5) the importance of reducing or eliminating use of fertilizers on lake front property; (6) the importance of minimizing transfer of AIS to the lake by having dedicated watercraft and cleaning boats that visit the lake.

Monitoring: Town Lakes Committee oversee(s) activity and assesses effectiveness.

Status: Anticipated to begin in 2020.



Recommended Actions for the Oxbow Lake APM Plan

Action #12: Identify and highlight high quality areas of littoral zone and riparian areas through review of aquatic plant and shoreland assessment data through various reports and online tools.

Objective: To (1) educate lake users on the value of these areas and the importance of good stewardship to their maintenance, (2) recognize landowners who implement good practices (e.g., large percentage of buffer area intact; three vegetative layers intact – herbaceous, shrubs, trees; areas of high native aquatic plant diversity and abundance), and (3) encourage landowners to implement good practices.

Monitoring: Town Lakes Committee and/or lake association promotes and oversees activity.

Status: Ongoing.

Action #13: Lake leaders should encourage and assist landowners to take on lake shore/shallow water improvement projects to rehabilitate areas identified through formal shoreland/shallow water assessments and/or lake user observations (sites might include areas of active erosion, channelized flow, point source pollution, impervious surfaces, and lawns) Vilas County Land and Water Conservation looks for partners in this endeavor and can provide planning and sponsorship of projects.

Objective: To rehabilitate specific areas of shoreland to improve natural functions and values.

Monitoring: Lake groups and lake leaders monitor and report progress to Town Lakes Committee.

Status: Ongoing.

Action #14: As part of an education program, encourage commitment from property owners to adopt practices that maintain/improve health of shoreland areas. In many cases, these are “practices” that mean less or no work (e.g., now mowing, no weed wacking, no leaf blowing, no removing large woody material).

Objective: To engage landowners in simple practices that improve/maintain health of the lake and shoreland.

Monitoring: Each landowner can monitor changes in the shoreland over time by simple means (e.g., annual mid-summer photographs or a catalog of plants and animals seen over time).

Status: Anticipated to begin in 2020.



Chapter 6. Contingency plan for AIS

Aquatic Invasive Species (AIS) are aquatic organisms that are introduced into waterbodies outside their natural range. These species include plants, animals, and pathogens (e.g., bacteria, fungi, and viruses). AIS may thrive in new environments because they lack environmental controls (such as natural predators). Waterbodies with AIS pose a risk to unaffected lakes through transportation of life stages by boats, trailers, and other vectors. Lake stewards should be prepared for the possibility of AIS colonization. Vigilant lake users are often the first to report an AIS in a waterbody. Early detection and response to an AIS can be helpful in limiting its success and spread.

For riparian owners and lake users, discovery of an AIS causes concern and an immediate desire to “fix the problem.” It is important to base responses to an AIS on scientific understanding of the particular species and the environment that it has colonized. A deliberate and systematic approach is beneficial. In some cases, the only management approach is to monitor how the lake ecosystem responds. For some AIS there is no effective treatment. For some AIS (especially plants) management is helpful, but careful consideration of the approach is essential since undesired outcomes of actions are problematic. An aquatic plant management plan (one including a contingency plan for AIS) is the best tool by which the process can be navigated. In fact, the APM plan is a requirement in Wisconsin for some kinds of aquatic plant management actions. One of the actions outlined in the previous chapter was to establish an Aquatic Invasive Species Rapid Response Team. This team and its coordinator are integral to the management process. It is important for this team to be multi-dimensional (or at least have quick access to the expertise that may be required). Since an AIS in a waterbody forms a source for colonization of other waterbodies, it is strategic for the Rapid Response Team to include representation from regional stakeholders.

Once the Rapid Response Team is organized, one of its first tasks is to develop a list of contacts and associated contact information (phone numbers and email addresses). At a minimum, this contact list should include: the Rapid Response Coordinator, members of the Rapid Response Team, County AIS Coordinator, WDNR Lakes Management Coordinator, Lake Association Presidents (or other points of contact), local WDNR warden, local government official(s), other experts, and contractor(s)/consultant(s). The Rapid Response Team should be prepared to respond to the possibility of any AIS (plant or animal). Since this contingency plan is part of an aquatic plant management plan, the remainder of this chapter focuses on aquatic invasive plant species.



Figure 5 provides a flowchart outlining an appropriate rapid response to the suspected discovery of an aquatic invasive plant species. The response will be most efficient if an AIS Rapid Response Team has already been established and is familiar with the contingency plan.

When a suspect aquatic invasive plant species is found, either the observer or a member of the Rapid Response Team (likely the coordinator) should collect an entire specimen including roots, stems, and flowers (if present). The sample should be placed in a sealable plastic bag with a small amount of water. Place a label in the bag written in pencil with date, time, collector's name, lake name, town, and county. Attach a lake map to the bag that marks the specific location of the suspect AIS and includes GPS coordinates (if available). The plant specimen should be placed in a refrigerator or on ice in a cooler. Deliver the sample to the WDNR Water Resource Management Specialist in the affected county as soon as possible (within three days). The WDNR or their botanical expert(s) will determine the species and confirm whether or not it is an AIS.

If the suspect plant specimen is judged to be an AIS, the next step is to determine the extent and density of the population since this will influence the management response. The Rapid Response Team should conduct (or have its consultant conduct) a survey to understand the distribution of the plant and, if appropriate, determine the perimeter and density for any beds. For rooted aquatic plant species, a "Pioneering Population" is designated when an invasive plant species covers a small area and is typically sparse and has been verified during the preceding 5 years. A "Pioneering Population" will cover an area that is less than three acres in size or has colonized less than 3% of the habitable area of the lake, stream reach, or wetland, whichever is greater. If greater than three acres (or >3% of the lake surface area) then it is designated an "Established Population." Once the infestation is characterized, "at risk" areas should also be marked on a map. For example, nearby boat landing sites and areas of high boat traffic.

When "pioneer" or "established" status is determined, it is time to consult the WDNR Lakes Coordinator to determine appropriate notifications and management responses. Necessary notifications of landowners, governmental officials, and recreationists (at boat landings) will be determined. Whether the population's perimeter needs to be marked with buoys will be decided by the WDNR. Funding sources will be identified and consultants and contractors will be contacted where necessary. The WDNR will determine if a further baseline plant survey is required (depending on type of treatment). A post treatment monitoring plan will be discussed and established to determine the efficacy of the selected treatment.






Figure 5. Aquatic Invasive Plant Species Rapid Response Flow Chart



Appendix 1. Literature cited



Hauxwell, J., S. Knight, K. Wagner, A. Mikulyuk, M. Nault, M. Porzky and S. Chase. 2010. *Recommended baseline monitoring of aquatic plants in Wisconsin: sampling design, field and laboratory procedures, data entry and analysis, and applications*. Wisconsin. Dept of Nat. Res. Bureau of Science Services, PUB-SS-1068 2010. Madison, WI.

Natural Heritage Inventory. 2021. *Elements by TownRange for Vilas County*. Retrieved 2021. http://dnr.wi.gov/topic/NHI/documents/Vilas_County.pdf

Nichols, Stanley A. 1999. *Floristic Quality Assessment of Wisconsin Lake Plant Communities with Example Applications*. Journal of Lake and Reservoir Management 15(2): 133-141.

Premo, Dean, Angie Stine, and Kent Premo. 2020. *Presque Isle Wilderness Waters Program: Oxbow Lake Aquatic Plant Management Plan*. White Water Associates, Inc.

Stine, Angie, Dean Premo, and Kent Premo. 2022. *The Wilderness Waters Adaptive Management Plan (Presque Isle Township, Vilas County, Wisconsin) - The Wilderness Waters Program*. White Water Associates, Inc.

US Department of Agriculture, Natural Resources Conservation Service. June 1986. *Urban Hydrology for Small Watersheds*. Technical Release-55.

Walters, C. 1986. *Objectives, constraints, and problem bounding*. In W.M. Getz, ed., *Adaptive Management of Renewable Resources*. Macmillan Publishing Company. New York, NY. p. 13+.

Wisconsin Administrative Code NR-102. 2024. *Water quality standards for Wisconsin surface waters*. Retrieved 2025. [WATER QUALITY STANDARDS FOR WISCONSIN SURFACE WATERS](#)

Wisconsin Department of Natural Resources. 2014. *Aquatic Invasive Species Early Detection Monitoring Standard Operating Procedure*. Retrieved 2017.

<http://dnr.wi.gov/water/wsSWIMSDocument.ashx?documentSeqNo=99459630>

Wisconsin Department of Natural Resources. July 20, 2020. *Lake Shoreland & Shallows Habitat Monitoring Field Protocol*. Wisconsin Department of Natural Resources.



Appendix 2. Documents



Appendix 2 - Table 1. Description of the aquatic plant community statistical terms

| Statistical term | Description |
|---|---|
| Maximum depth of plants (ft) | Since aquatic rooted aquatic plants require sunlight at the lake bottom in order to root and grow, water depth and transparency ultimately limit where plants can grow. By repeated sampling in a lake, we can determine the maximum depth at which plants grow. By definition this area where plants can grow is called the littoral zone. Beyond that depth plants are absent. |
| Frequency of occurrence at points within the littoral zone | Not every area of the lake bottom that is shallower than the maximum depth of plants actually has plants living there. Other things like the kind of substrate can influence whether a plant can take root. Frequency of occurrence at sites shallower than maximum depth of plants documents the proportion of sites where rooted plants actually occur within the littoral zone. |
| Average number of all species per point (vegetative. points only) | This metric is the average of the number of plant species documented at each of the point-intercept sampling points. Lakes with diverse plant communities tend to have higher average number of plant species at each site. |
| Species richness | This metric is a simple count of the number of species in an area. In the case of the point-intercept survey, this refers to the number of plant species actually collected on the rake for the point-intercept sampling points in a given area (e.g., an entire lake or a sub-PI plot). The species richness of a lake would mean the total number of plant species in the entire lake. Species richness is a simple measure of diversity. More species equals higher diversity. |
| Species richness (including visuals) | This metric includes the simple count of the number of plant species collected on the rake for the point-intercept sampling points in a given area and those observed (but not actually collected with the rake) within 6 feet of the sampling points. |
| Simpson diversity Index | This is a more sophisticated measure of biological diversity than species richness. It is an index that takes into account species richness (number of species) as well as evenness. Species evenness is a description of the distribution of abundance across the species in a community. Species evenness is highest when all species in a sample have the same abundance. The most diverse systems have both high number of species and high evenness. |



| Statistical term | Description |
|-------------------------------|--|
| Floristic Quality Index (FQI) | This is an index that tries to capture how “pristine” a plant community is in terms of its species composition. Some plant species are very sensitive to human presence and are less likely to occur in lakes where human influences are strong. At the other end of the spectrum are plants that do very well in the presence of human influence. The FQI attempts to evaluate the mix of sensitive and non-sensitive plants in a lake. |
| Average rake fullness | Rake fullness is a rating given to each rake pull on a PI survey that indicates the amount of plant material for each species as well as combined species. There are four ranks: 0 (not present), 1, 2, & 3 (3 represents the highest value of plant material on a rake). Average rake fullness is the mean value for all the point-intercept points. |



Appendix 2 - Table 2. Plant species recorded and statistics for the 2024 Oxbow Lake aquatic plant survey.

| Common name | Scientific name | Frequency of occurrence within vegetated areas (%) | Frequency of occurrence at sites shallower than maximum depth of plants | Relative Frequency (%) | Number of sites where species found (including visuals) | Average Rake Fullness |
|--------------------------|---|--|---|------------------------|---|-----------------------|
| Slender naiad | <i>Najas flexilis</i> | 41.78 | 24.48 | 17.41 | 96 | 1.11 |
| Watershield | <i>Brasenia schreberi</i> | 24.00 | 14.06 | 10.00 | 90 | 1.41 |
| Large-leaf pondweed | <i>Potamogeton amplifolius</i> | 21.33 | 12.50 | 8.89 | 102 | 1.15 |
| Muskgrasses | <i>Chara sp.</i> | 20.44 | 11.98 | 8.52 | 48 | 1.39 |
| Berchtold's pondweed | <i>Potamogeton berchtoldii</i> | 20.00 | 11.72 | 8.33 | 47 | 1.11 |
| Northern naiad | <i>Najas gracillima</i> | 18.22 | 10.68 | 7.59 | 42 | 1.12 |
| Nitella | <i>Nitella sp.</i> | 12.44 | 7.29 | 5.19 | 29 | 1.00 |
| Large purple bladderwort | <i>Utricularia purpurea</i> | 10.22 | 5.99 | 4.26 | 32 | 1.00 |
| Creeping spikerush | <i>Eleocharis palustris</i> | 8.00 | 4.69 | 3.33 | 42 | 1.06 |
| Variable pondweed | <i>Potamogeton gramineus</i> | 8.00 | 4.69 | 3.33 | 36 | 1.28 |
| Freshwater sponge | <i>Freshwater sponge</i> | 7.56 | 4.43 | | 22 | 1.00 |
| Water marigold | <i>Bidens beckii</i> (formerly <i>Megalodonta</i>) | 5.78 | 3.39 | 2.41 | 15 | 1.00 |
| Slender waterweed | <i>Elodea nuttallii</i> | 5.78 | 3.39 | 2.41 | 14 | 1.15 |



| Common name | Scientific name | Frequency of occurrence within vegetated areas (%) | Frequency of occurrence at sites shallower than maximum depth of plants | Relative Frequency (%) | Number of sites where species found (including visuals) | Average Rake Fullness |
|--------------------------|---------------------------------|--|---|------------------------|---|-----------------------|
| Floating-leaf bur-reed | <i>Sparganium fluctuans</i> | 5.33 | 3.13 | 2.22 | 19 | 1.25 |
| Spiral-fruited pondweed | <i>Potamogeton spirillus</i> | 4.44 | 2.60 | 1.85 | 13 | 1.00 |
| Needle spikerush | <i>Eleocharis acicularis</i> | 4.00 | 2.34 | 1.67 | 13 | 1.00 |
| Wild celery | <i>Vallisneria americana</i> | 3.56 | 2.08 | 1.48 | 11 | 1.00 |
| Fern pondweed | <i>Potamogeton robbinsii</i> | 3.11 | 1.82 | 1.30 | 7 | 1.00 |
| Vasey's pondweed | <i>Potamogeton vaseyi</i> | 2.67 | 1.56 | 1.11 | 6 | 1.00 |
| Spiny hornwort | <i>Ceratophyllum echinatum</i> | 2.22 | 1.30 | 0.93 | 5 | 1.00 |
| Ribbon-leaf pondweed | <i>Potamogeton epihydrus</i> | 2.22 | 1.30 | 0.93 | 29 | 1.00 |
| Common waterweed | <i>Elodea canadensis</i> | 2.22 | 1.30 | 0.93 | 6 | 1.00 |
| Pickerelweed | <i>Pontederia cordata</i> | 1.78 | 1.04 | 0.74 | 25 | 1.00 |
| Spatterdock | <i>Nuphar variegata</i> | 1.78 | 1.04 | 0.74 | 15 | 1.00 |
| Blunt-leaf pondweed | <i>Potamogeton obtusifolius</i> | 1.33 | 0.78 | 0.56 | 3 | 1.00 |
| Small purple bladderwort | <i>Utricularia resupinata</i> | 1.33 | 0.78 | 0.56 | 3 | 1.00 |
| Water horsetail | <i>Equisetum fluviatile</i> | 0.89 | 0.52 | 0.37 | 7 | 1.00 |
| Grass-leaved arrowhead | <i>Sagittaria graminea</i> | 0.89 | 0.52 | 0.37 | 7 | 1.00 |




| Common name | Scientific name | Frequency of occurrence within vegetated areas (%) | Frequency of occurrence at sites shallower than maximum depth of plants | Relative Frequency (%) | Number of sites where species found (including visuals) | Average Rake Fullness |
|-------------------------|---------------------------------------|--|---|------------------------|---|-----------------------|
| Quillwort | <i>Isoetes sp.</i> | 0.89 | 0.52 | 0.37 | 6 | 1.00 |
| Brown-fruited rush | <i>Juncus pelocarpus f. submersus</i> | 0.89 | 0.52 | 0.37 | 4 | 1.00 |
| Twig rush | <i>Cladium mariscoides</i> | 0.89 | 0.52 | 0.37 | 4 | 1.00 |
| Leafy pondweed | <i>Potamogeton foliosus</i> | 0.44 | 0.26 | 0.19 | 1 | 1.00 |
| Hardstem bulrush | <i>Schoenoplectus acutus</i> | 0.44 | 0.26 | 0.19 | 1 | 1.00 |
| Narrow-leaved bur-reed | <i>Sparganium angustifolium</i> | 0.44 | 0.26 | 0.19 | 1 | 1.00 |
| Aquatic moss | <i>Aquatic moss</i> | 0.44 | 0.26 | | 1 | 1.00 |
| Filamentous algae | <i>Filamentous algae</i> | 0.44 | 0.26 | | 1 | 1.00 |
| White water lily | <i>Nymphaea odorata</i> | 0.44 | 0.26 | 0.19 | 12 | 1.00 |
| Three-way sedge | <i>Dulichium arundinaceum</i> | 0.44 | 0.26 | 0.19 | 10 | 1.00 |
| Water lobelia | <i>Lobelia dortmanna</i> | 0.44 | 0.26 | 0.19 | 6 | 1.00 |
| Common bladderwort | <i>Utricularia vulgaris</i> | 0.44 | 0.26 | 0.19 | 4 | 1.00 |
| Pipewort | <i>Eriocaulon aquaticum</i> | 0.44 | 0.26 | 0.19 | 3 | 1.00 |
| Arrowhead | <i>Sagittaria sp.</i> | | | | 2 | |
| Farwell's water-milfoil | <i>Myriophyllum farwellii</i> | | | | 1 | |




| Common name | Scientific name | Frequency of occurrence within vegetated areas (%) | Frequency of occurrence at sites shallower than maximum depth of plants | Relative Frequency (%) | Number of sites where species found (including visuals) | Average Rake Fullness |
|---------------------|------------------------------|--|---|------------------------|---|-----------------------|
| Dwarf water-milfoil | <i>Myriophyllum tenellum</i> | | | | 1 | |
| Iris sp. | <i>Iris sp.</i> | | | | 1 | |






Appendix 2 - Map 1. North Part of Oxbow Lake total rake fullness distribution during the 2014 aquatic plant survey.






Appendix 2 - Map 2. North Part of Oxbow Lake total rake fullness distribution during the 2018 aquatic plant survey.






Appendix 2 - Map 3. North Part of Oxbow Lake total rake fullness distribution during the 2024 aquatic plant survey.






Appendix 2 - Map 4. South Part of Oxbow Lake total rake fullness distribution during the 2014 aquatic plant survey.





Appendix 2 - Map 5. South Part of Oxbow Lake total rake fullness distribution during the 2018 aquatic plant survey.





Appendix 2 - Map 6. South Part of Oxbow Lake total rake fullness distribution during the 2024 aquatic plant survey.



Appendix 3. Aquatic Invasive Species Report



Oxbow Lake (Vilas County, Wisconsin) Aquatic Invasive Species Monitoring Report

Prepared for:

Presque Isle Town Lake Committee
Contact: Nick Williams
P.O. Box 37
Presque Isle, Wisconsin 54557
Phone: (715) 892-7184; E-mail: thewds@milwpc.com

Prepared by:

White Water Associates, Inc.
Contact: Carole Gabbard
429 River Lane, P.O. Box 27
Amasa, Michigan 49903
Phone: (906) 822-7889; Email: carole.gabbard@white-water-associates.com

Cite as: *Gabbard, Carole. 2024. Aquatic Invasive Species Monitoring Report for Oxbow Lake (Vilas County, Wisconsin). White Water Associates, Inc.*



Date: December 2025

INTRODUCTION

White Water Associates, Inc. (hereafter referred to as White Water) was retained by the Presque Isle Town Lake Committee to conduct an Aquatic Invasive Species (AIS) monitoring in Oxbow Lake (Vilas County, Wisconsin) associated with an aquatic plant survey in 2024.


METHODS

For this survey, we conducted a thorough search of likely habitats for aquatic invasive species (AIS) around the entire Oxbow Lake shoreline with special focus on the boat landing and other points of most likely AIS introduction or colonization. The field work was conducted on August 20, 2024 by White Water aquatic biologists Carole Gabbard and Jesse Gabbard and field technician Faryn Rice. The late season timing was to target aquatic invasive species that are more easily identified in the late season (for example, purple loosestrife and narrow-leaved cattail).

The search began at the public boat landing. Using the boat, we meander-searched around the entire lake and stopped at five target search sites and snorkeled and waded with viewing bucket to look for AIS in the water and near shore. As part of the survey, a Ponar dredge was used to collect a sediment sample at the deep hole location in the lake. This sample was later examined in the laboratory using a stereo microscope to look for spiny water flea body parts. Spiny water fleas (*Bythotrephes longimanus*) are an aquatic invasive zooplankton that is found in a few lakes in Wisconsin. The survey ended by wading and snorkeling at the boat landing area looking for AIS. The shoreline in the vicinity of the boat landing was walked and waded looking for AIS (Map 1).



Oxbow Lake - 2024 Aquatic Invasive Species Report



Map 1. Oxbow Lake target search sites (OX2024-#), dredge (OX2024-DR) and boat landing locations.



SURVEY RESULTS

One wetland invasive plant species was observed: aquatic forget-me-not (*Myosotis scorpioides*). During this survey, vegetative parts of an Iris species were observed. This plant could be the wetland invasive species known as yellow iris (*Iris pseudacorus*). Each of these invasive species is described in the following paragraphs. The spiny water flea (*Bythotrephes longimanus*) was not found in the Oxbow Lake sediment sample.

1. **Aquatic forget-me-not** (Photo 1). This invasive species can quickly crowd native plant species and is able to form large monocultures. This, in turn, affects community compositions by reducing the number of native herbs (Wisconsin DNR, 2025). Aquatic forget-me-not is difficult to control due to its mechanisms for spreading (Wisconsin DNR, 2025). Aquatic forget-me-not was found at one location during the 2024 AIS survey (Map 2).
2. **Yellow iris**. At the time of the 2024 survey, iris plants were not in flower making definitive species identification impossible (Photo 2). The locations of the iris found are shown in the Map 2. We recommend monitoring the population annually in the spring (June) to confirm the presence of yellow iris by the blossom color. If yellow iris plants are present, they should be removed by digging up the entire plant (including the underground rhizome). To stabilize the resulting disturbed shoreline, we recommend that native blue flag iris (*Iris versicolor*) be planted. This species is available from nurseries.

While conducting the aquatic plant and aquatic invasive species survey, we observed native phragmites (American common reed, *Phragmites australis subsp. Americanus*) at one location on Oxbow Lake (GPS coordinates 46.24326, - 89.67392). We collected and pressed specimens and provided them to Robert W. Freckmann (UW-Stevens Point Herbarium) who confirmed our identification. You can find online several websites to help identify the native to the non-native phragmites. Below are some examples of useful information online:

[Identifying Native vs. Invasive Phragmites | Great Lakes Phragmites Collaborative](#)
[Invasive-Phragmites - University of Wisconsin-Madison.pdf](#)





Photo 1. Aquatic forget-me-not (tiny blue flowers) on a northern Wisconsin Lake shoreline.



Photo 2. Possible yellow iris found on a northern Wisconsin Lake shoreline.




Oxbow Lake - 2024 Aquatic Invasive Species Report

Table 1. Target search site GPS coordinates and their findings.

| Target search site | Latitude | Longitude | Findings |
|---------------------------|-----------------|------------------|----------------------|
| OX2024-01 | 46.22696 | -89.68419 | Possible yellow iris |
| OX2024-02 | 46.24755 | -89.66438 | No AIS Present |
| OX2024-03 | 46.23908 | -89.68624 | No AIS Present |
| OX2024-04 | 46.24908 | -89.68704 | No AIS Present |
| OX2024-05 | 46.23538 | -89.68036 | No AIS Present |
| Boat Landing | 46.23721 | -89.67422 | No AIS Present |
| OX2024-DR | 46.24626 | -89.68811 | No AIS Present |





Map 2. Wetland invasive species found during the 2024 AIS in Oxbow Lake.

Literature cited

Wisconsin DNR, 2025. [Aquatic Forget-me-not | \(Myosotis scorpioides\) | Wisconsin DNR](#)

